

On the spin asymmetry of ground states in trapped two-component Fermi gases with repulsive interactions

M. Ögren, K. Kärkkäinen, Y. Yu and S.M. Reimann

Mathematical Physics, Lund Institute of Technology, P.O. Box 118, SE-22100 Lund, Sweden

(Dated: February 6, 2008)

We examine the spin asymmetry of ground states for two-dimensional, harmonically trapped two-component gases of fermionic atoms at zero temperature with weakly repulsive short range interactions. Our main result is that, in contrast to the three-dimensional case, in two dimensions a non-trivial spin-asymmetric phase can only be caused by shell structure. A simple, qualitative description is given in terms of an approximate single particle model, comparing well to the standard results of Hartree-Fock or direct diagonalization methods.

PACS numbers: 05.30Fk, 03.65.Ge

I. INTRODUCTION

Cold atomic gases with multiple components have been realized for example in Bose-Einstein condensates [1, 2, 3, 4]. The nowadays possible experimental studies of atomic Fermi gases [5, 6, 7, 8, 9, 10, 11, 12, 13] have raised much recent interest in the different underlying quantum phases, with a main attention on pairing. Fermi gases with spin imbalance have been a hot topic more recently in connection with superfluidity issues [14]. While most of the previous studies focus on attractive interactions, Duine and MacDonald [15] pointed out that the repulsive regime might offer the possibility of ferromagnetism in ultra-cold atom gases with, as they say, ‘unprecedented experimental control’. Describing spin (magnetic) properties of Fermi systems is an important issue in many-body physics. The trapped dilute fermionic atom gases, with their flexibility in tuning important system parameters such as the interaction strength, could provide opportunities to look for new magnetic phenomena and to test the theoretical models of magnetic properties.

Possible spin symmetry breaking for a Fermi gas in the unitarity limit [16, 17, 18, 19, 20] was discussed by Chevy [21]. Maruyama and Bertsch [22] have included time-dependent, external magnetic fields, yielding spin-asymmetric ground states.

Sogo and Yabu [23] studied the three-dimensional, trapped Fermi gas with two components. (The more general multi-component problem was discussed earlier by Salasnich *et al.* [24]). These components can be characterized by different intrinsic spin states, or for example, two hyperfine states, playing the role of pseudospin in a two-component Fermi gas. For weak interactions between the components, the zero-temperature ground state is symmetric, with equal up- and down-densities and vanishing total spin or pseudospin. Sogo and Yabu showed, that in three spatial dimensions, depending on the interaction strength between the components, a spin-asymmetric, so-called “collective ferromagnetic” state may become energetically favorable. This spin or pseudospin asymmetry in the ground state arises

as it becomes energetically favorable to increase the Fermi energy of one component rather than increasing the interaction energy between the two species. Sogo and Yabu [23] based their investigation on the Thomas-Fermi approximation [25, 26], an efficient way to compute the ground state properties of a large system – however, at the cost of ignoring the shell corrections [25].

These studies were performed for a three-dimensional harmonic trap. However, by appropriate manipulation of the trapping potential, confined ultra-cold atoms can achieve conditions of reduced dimensionality. We thus found it useful to further investigate the question of spin asymmetry for an isotropic harmonic trap in *two dimensions*. (In one dimension, a uniform fermi system can be solved exactly with a Bethe ansatz [27]. Recently, this was also generalized to a harmonic trap within the local density approximation [28]).

We found that in two dimensions the spin asymmetry can only arise from the shell-structure and Hund’s rule.

II. THE THOMAS-FERMI APPROXIMATION FOR A TWO-COMPONENT FERMION GAS

Consider fermionic particles of equal mass m at zero temperature, with two internal (spin) states labeled by $\sigma \in \{u, d\}$. The fermions shall be subject to repulsive two-body contact interactions. In atomic units ($\hbar = \omega = m = 1$), our model Hamiltonian in two dimensions is ($r = |\mathbf{r}|$)

$$H = \sum_{\sigma} \sum_i^{N_{\sigma}} \left[-\frac{1}{2} \Delta_i + \frac{1}{2} r_i^2 \right] + g \sum_{i,u < j,d} \delta(\mathbf{r}_i - \mathbf{r}_j), \quad (1)$$

where N_{σ} denotes the number of particles in each of the two spin states. Here, only the repulsive, weak-interaction regime of Eq. (1) is considered. The coupling constant $g \propto a/l_z > 0$ is proportional to the ratio of the three-dimensional scattering length a and the length scale l_z of the ground state in the frozen-out z -dimension. From the condition that the (mean-field)

interaction energy should be much smaller than the Fermi energy μ_F

$$gn_u^0/\mu_F \ll 1, \quad (2)$$

with n_u^0 denoting the particle density (of one of the species) in the center of the trap, one is led to the diluteness condition

$$0 \leq g \ll 2\pi. \quad (3)$$

We start by treating the problem within the Thomas Fermi (TF) approximation which should be valid in the limit of large particle numbers where the specific contribution from shell structure plays a minor role. In the Thomas-Fermi approximation, the total energy is

$$E = \int d\mathbf{r} \left[\pi (n_u^2 + n_d^2) + gn_u n_d + \frac{1}{2} r^2 n \right], \quad (4)$$

where n_u and n_d and $n = n_u + n_d$ are the (pseudo)spin- and total densities, respectively. The Euler-Lagrange equations for the densities, while fulfilling the constraint of constant particle number $N = N_u + N_d = \int d\mathbf{r} n$, are

$$\frac{\delta(E - \mu_\sigma N_\sigma)}{\delta n_\sigma} = 0 \Rightarrow \begin{cases} 2\pi n_u + gn_d + r^2/2 = \mu_u \\ 2\pi n_d + gn_u + r^2/2 = \mu_d \end{cases} \quad (5)$$

In order for the system to be in equilibrium we must have $\mu_u = \mu_d \equiv \mu$. Further on for the system (5) to be a stable solution we must have

$$\left| \frac{\frac{\delta^2 E}{\delta n_u^2}}{\frac{\delta^2 E}{\delta n_d \delta n_u}} \quad \frac{\frac{\delta^2 E}{\delta n_u \delta n_d}}{\frac{\delta^2 E}{\delta n_d^2}} \right| \geq 0 \Leftrightarrow g \leq 2\pi, \quad (6)$$

which is guaranteed by the diluteness condition. By viewing (5) as a matrix equation in the variables n_u and n_d it has the system determinant $4\pi^2 - g^2$. So for $g < 2\pi$, which is again guaranteed by the diluteness condition, the system (5) has a unique algebraic solution which corresponds to a stable minimizer of Eq. (4)

$$\begin{cases} n_u = \frac{1}{2\pi+g} \left(\mu - \frac{1}{2} r^2 \right), & r \leq \sqrt{2\mu} \\ n_d = n_u \end{cases} \quad (7)$$

This TF result suggests that there should be no stable spin-asymmetric density configurations in the two-dimensional system under study. This is qualitatively different from the corresponding TF investigation for three-dimensional system where a non-trivial phase separation called “collective ferromagnetic state” has been reported [23, 24]. This is due to the different exponent of the kinetic energy functional in the three-dimensional case, leading to non-linear equations for the densities.

Inserting the density (7), $n = n_u + n_d = 2n_u$, into the energy functional (4), the total energy equals

$$E = \frac{2}{3} \sqrt{1 + \frac{g}{2\pi}} N^{3/2}. \quad (8)$$

We then note that for $g = 2\pi$ this energy coincides with the fully polarized phase, i.e. 1/2 times Eq. (8) with $g = 0$ and $N \rightarrow 2N$, hence giving a physical motivation for the condition in Eq. (6). This describes the trivial phase separation to a totally polarized system similar to what can happen in three dimensions [23].

III. THOMAS-FERMI SINGLE-PARTICLE MODEL

Let us now incorporate the shell structure. Obviously, the density profile given by the above Eq. (7) will also be valid for two unequally populated non-interacting ($g = 0$) spin states, as according to Eq. (5), n_u and n_d are then uncoupled. This is the situation when particles of one (pseudo)spin state are filling a higher shell than the other spin state, such that $|\mu_u - \mu_d| = \omega_{eff}$ (ω_{eff} is defined in Eq. (11)). The major effect of the repulsive interaction is to redistribute atoms from the center of the cloud to larger r values. This is qualitatively modeled by the profile of the upper Eq. (7) also when $|\mu_u - \mu_d| = \omega_{eff}$ (and agrees well with a numerical solution of Eq. (5) for $g \ll 2\pi$). Moreover, the profile of Eq. (7) does not break the $SU(2)$ symmetry, as also was shown to be the case for the exact density [29].

We now begin from the single-particle Hartree-Fock equations [25] for the Hamiltonian (1) in the weakly repulsive regime

$$\left[-\frac{1}{2} \Delta + gn_d(x, y) + \frac{1}{2} (x^2 + y^2) \right] \varphi_{i,u} = E_{i,u} \varphi_{i,u}. \quad (9)$$

Now, for the purpose of presenting a simple analytical model, we approximate the particle density with the g -dependent TF density corresponding to Eq. (7)

$$n_d(x, y) \approx n_d^{TF}(|\mathbf{r}|) \equiv$$

$$\frac{1}{2\pi + g} \left(\mu_d - \frac{1}{2} (x^2 + y^2) \right) \theta \left(\mu_d - \frac{1}{2} (x^2 + y^2) \right),$$

where $\mu_d = \sqrt{(2 + g/\pi) N_d}$ and $\theta(\cdot)$ is Heaviside's step-function. After separating the wavefunctions $\varphi_{i,u}(x, y)$ in Eq. (9) into the Cartesian coordinates we obtain e.g. for the x -dimension

$$\left[-\frac{1}{2} \frac{d^2}{dx^2} + \underbrace{\frac{1}{2} \left(1 - \frac{g\theta(\mu_d - x^2/2)}{2\pi + g} \right)}_{\omega_{eff}^2(g, N_d)} x^2 \right] \psi_{i,u}(x) = \quad (10)$$

$$\left[\epsilon_{i,u} - \underbrace{\frac{g\mu_d\theta(\mu_d - x^2/2)}{2\pi + g}}_{\epsilon_{shift}(g, N_d)} \right] \psi_{i,u}(x). \quad (11)$$




Figure 1: (Color online) The N_u - N_d chart for harmonically trapped 2D fermions with two spin states according to TFSPM (here with $g = 0.1$). The black dots ($\bullet$) show the states with lowest energy for a given $N = N_u + N_d$, the corresponding spin for this states are seen in the inset figure. The green triangles ($\triangle$) show case *ii*) (*spin symmetry*) from equation (16), green circles ($\circ$) show case *iii*) (*polarized*). The red stars ($\star$) are the magic numbers $M_j = j(j+1) = 2, 6, 12, \dots$

By re-arranging the terms in Eq. (11), this can be viewed as a one-dimensional harmonic oscillator with the perturbation

$$h_p = \left(\frac{g\mu_d}{2\pi + g} - \frac{1}{2} \frac{g}{2\pi + g} x^2 \right) \theta(\mu_d - x^2/2). \quad (12)$$

We then apply first order non-degenerate perturbation theory in each Cartesian dimension

$$\epsilon_{i,u} \approx i + \frac{1}{2} + \langle \phi_i | h_p | \phi_i \rangle, \quad (13)$$

where ϕ_i is the eigenfunction $i = 0, 1, 2, \dots$ to the one-dimensional harmonic oscillator with frequency $\omega = 1$. If we now only keep the leading order terms in $g \ll 2\pi$, we can for practical purposes write Eq. (13) as

$$\epsilon_{i,u} \approx i + \frac{1}{2} + \frac{g}{2\pi} \int_{x^2 < \sqrt{8N_d}} \left(\sqrt{2N_d} - \frac{1}{2} x^2 \right) |\phi_i(x)|^2 dx. \quad (14)$$

Clearly this is the x -dependent part of what one obtains if one treats the mean-field interaction term in Eq. (9)

as gn_{HO} from the beginning, where n_{HO} refers to the TF-density of the non-interacting system.

Obviously, Eq. (13) takes the value for the non-interacting HO $\epsilon_{i,u} = i + 1/2$ in the two limits $g \rightarrow 0$ and $N_d \rightarrow 0$ ($N_u \rightarrow N$). In total, we have two Cartesian coordinates and two spin-components such that the total energy is [25]

$$E_{tot} = E_u + E_d \approx 2 \sum_{occ.} (\epsilon_{i,u} + \epsilon_{i,d}) - g \int d\mathbf{r} n_u n_d. \quad (15)$$

At zero temperature we have a Fermi-Dirac occupation of each component such that the HO main quantum number increases by one unit whenever a shell is filled. There are three special cases for which the total energy (15) can easily be written down explicitly.

i) The non-interacting case: When $g = 0$ we have $\epsilon_{i,\sigma} = i + 1/2$.

ii) Spin symmetry: When $N_u = N_d$, before subtracting the last term in Eq. (15), the total energy is simply twice the one for a single component system with the number of particles being $N/2$ and with a poten-

tial $V(g, N/2) = \omega_{\text{eff}}^2 r^2/2 + \epsilon_{\text{shift}}$, but in this case the Thomas-Fermi radius $R_{TF} = \sqrt{2\mu_{TF}}$ is equal for both components. Then all the single particle wavefunctions decay very rapidly for $r > R_{TF}$ and we can formally drop the factors $\theta(\mu_d - x^2/2)$ in Eq. (11), then $\epsilon_{i,\sigma} = \omega_{\text{eff}}(i + 1/2) + \epsilon_{\text{shift}}$.

iii) The totally polarized case: When $N = N_u$ and $N_d = 0$ (or vice versa) the interaction disappears which gives $\epsilon_{i,u} = i + 1/2$. The cases *i*), *ii*) and *iii*) are all captured by the formula [30]

$$E_{u,d}(g, \alpha) = \sqrt{\frac{2\pi}{2\pi + g}} \left[\frac{1}{6} \text{ceil} \left(\frac{\sqrt{1 + 8\alpha} - 1}{2} \right) \times \left(6\alpha + 1 - \left(\text{ceil} \left(\frac{\sqrt{1 + 8\alpha} - 1}{2} \right) \right)^2 \right) + \frac{g}{\sqrt{2\pi}} \alpha^{3/2} \right], \quad (16)$$

where $\text{ceil}(\cdot)$ is the closest integer from above. For case *i*) one has $E_{\text{tot}} = E_u(g=0, \alpha=N_u) + E_d(g=0, \alpha=N_d)$. In case *ii*) $E_{\text{tot}} = 2E_{u,d}(g, \alpha=N/2) - e$ where $e = g \int d\mathbf{r} n^2/4 = gN^{3/2}/\sqrt{36\pi^2 + 18\pi g}$. In case *iii*) $E_{\text{tot}} = E_{u,d}(g=0, \alpha=N)$.

For all other cases, Eq. (14) can be calculated (numerically) for each single particle level $\epsilon_{i,\sigma}$ which are then summed up according to Eq. (15). If one compares this procedure with a case where Eq. (16) is applicable e.g. case *ii*) where $N_d = N_u = N/2$, one finds that the relative difference between those two approaches is small (e.g. $< 2 \cdot 10^{-3}$ for $g = 0.1$ and $N = 10$) and decreases with N .

The approximative model we suggest here to illustrate shell-induced spin asymmetry relies upon the fact that a spherically symmetric TF particle-density $n_{TF} \propto (\mu - r^2/2)$ does not break the $SU(2)$ symmetry, which is present for the non-interacting problem due to the harmonic trap. It has been shown in lowest order perturbation theory that the ℓ -degeneracy for a given HO shell is not lifted, even when the exact expression for the particle-density is used [29]. This fact supports the validity of our approximative “Thomas-Fermi single particle model”, TFSPM, for weak interactions. However, except for the noninteracting case ($g = 0$), the TFSPM obviously overestimates the many-body energy compared to more sophisticated computational methods, where there is more freedom to vary the many-body wavefunction. TFSPM successfully finds the correct spin-configurations with lowest energy, see Figure 1, and reproduces the shell correction with very high accuracy, see Figure 2. What we here refer to as the correct spin-configuration is what we obtained from full Hartree-Fock calculations (see below) for all presented values of N . For $N \leq 12$ we confirmed the same spin-configuration by direct numerical diagonalization of the Hamiltonian (1).

IV. COMPARISON TO THE HARTREE-FOCK METHOD

Now we turn our interest to the full Hartree-Fock calculation. We calculate the particle densities (e.g.) $n_d(x, y) = \sum_{i=1}^{N_d} |\varphi_{i,d}|^2$ (without demanding spherical symmetry) for each spin component iteratively from the mean-field single-particle states $\varphi_{i,d}(x, y)$ corresponding to the Hartree-Fock Eq. (9).

For each total number of particles $N = N_u + N_d$, we determine the lowest-energy state self-consistently. For this state, the total spin is defined as $S = (N_u - N_d)/2$. A zig-zag pattern with increasing amplitude and frequency occurs, see inset of Figure 1, which is identical for the Hartree-Fock calculations and the TFSPM. This pattern (inset of Figure 1) is a direct consequence of Hund’s rule. The trapped atoms at a degenerate shell can lower their interaction energy by maximizing the number of atoms of the same species - in other words, by aligning their spins. This mechanism in contact-interacting fermionic systems leads to Hund’s first rule and what we here call shell-induced spin asymmetry, in close similarity to long-range interacting systems. Here, however, Hund’s rule has a more dramatic effect since it removes completely the interaction between atoms of the same spin state.

However, in absolute values, the energy difference associated with this pattern is a decreasing function of the number of particles. To be more specific, the lowest energy state carrying the largest total spin in each shell is $S_{\text{max}}(j) = (j + 1)/2$, where $j = 1, 2, 3, \dots$ is the index of the shell. Since the magic numbers of an isotropic two-dimensional harmonic oscillator with two spin-states are $M_j = 2 \sum_{m=0}^{j-1} (m + 1) = j(j + 1) = 2, 6, 12, \dots$, S_{max} is related to the total number of particles as $S_{\text{max}} \simeq \sqrt{N}/2$. The energy difference of the state with $S(N) = S_{\text{max}}$ compared to the state $S(N) = 0$ is of the order $\Delta E \sim g\sqrt{N}$. Comparing this to the leading order term in the TF estimate of the total energy of the state with $S(N) = 0$ from Eq. (8) $E \simeq 2N^{3/2}/3$, the relative energy gain of the shell-induced spin asymmetry scales as $\Delta E/E \sim g/N$, which becomes irrelevant for large N . This is in agreement with the analysis in section II, which had shown that the Thomas-Fermi approximation could not predict any stable spin-asymmetric phase for the system, but only the trivial, totally polarised system.

V. CONCLUSIONS

In summary, the short-range weakly repulsive interacting fermi gas is a simple toy model that illustrates very directly Hund’s rule: The total spin is maximized at mid-shell, in order to minimize the total energy, what we here have referred to as shell-induced spin asymmetry. For the case of an isotropic two-dimensional harmonic trap we have illustrated a simple approximative single




Figure 2: (Color online) Oscillating part of the total energy (shell correction) for the state with lowest energy, at a given N (here with $g = 0.1$). The solid black line with stars ($\star$) shows the TFSPM, the red dashed line with circles ($\circ$) the numerical Hartree-Fock results. The green triangles ($\triangle$) show case *ii*) (*spin symmetry*) from equation (16), which is only defined for even N . In this case the total energy was $\sim 0.3\%$ lower for Hartree-Fock compared to TFSPM and the deviations in the shell correction are very small. Investigating even smaller values of g the curves approach each other further. The smooth part of the energy is constructed such that the different curves for the oscillating part should coincide at the minima of the shell correction (i.e. at the magic numbers) and they follow a line $y \propto N^{1/2}$.

particle model (TFSPM), that qualitatively reproduces the energy landscape of the system, when compared to Hartree-Fock calculations. This model is restricted to two dimensions and weak interactions since it relies upon the fact that the TF density profile does not break the $SU(2)$ symmetry of the non-interacting problem and the fact that perturbation theory is used. This also outrules the possibility to see super-shell structure, reported in the three-dimensional case [31], within this approach.

ACKNOWLEDGEMENTS

This work was financially supported by the Swedish Research Council and the Swedish Foundation for Strategic Research, as well as the European Community project ULTRA-1D (NMP4-CT-2003-505457).

[1] M.H. Anderson, J.R. Ensher, M. R. Matthews, C.E. Wieman and E.A. Cornell, *Science* **269**, 189 (1995).

[2] K.B. Davis, M.O. Mewes, M.R. Andrews, N.J. van Druten, D.S. Druffee, D.M. Kurn and W. Ketterle, *Phys. Rev. Lett.* **75**, 3969 (1995).

[3] W. Ketterle, *Rev. Mod. Phys.* **74**, 1131 (2002).

[4] E. A. Cornell and C. E. Wieman, *Rev. Mod. Phys.* **74**, 875 (2002).

[5] B. De Marco and D.S. Jin, *Science* **285**, 1703 (1999).

[6] M.J. Holland, B. DeMarco, and D.S. Jin, *Phys. Rev. A* **61**, 053610 (2000).

[7] S.R. Granade, N.E. Gehm, K.M. O'Hara, and J.E. Thomas, *Phys. Rev. Lett.* **88**, 120405 (2002).

[8] S. Jochim, M. Bartenstein, G. Hendl, J. Hecker Denchlag, and R. Grimm, *Phys. Rev. Lett.* **89**, 273202 (2002).

[9] C.A. Regal and D.S. Jin, *Phys. Rev. Lett.* **90**, 230404 (2003).

[10] Z. Hadzibabic, S. Gupta, C. A. Stan, C. H. Schunck, M. W. Zwierlein, K. Dieckmann, and W. Ketterle, *Phys. Rev. Lett.* **91**, 160401 (2003).

[11] M. Greiner, C.A. Regal and D.S. Jin, *Nature (London)* **426**, 537 (2003).

[12] M. W. Zwierlein, C. A. Stan, C. H. Schunck, S. M. F. Raupach, S. Gupta, Z. Hadzibabic, and W. Ketterle, *Phys. Rev. Lett.* **91**, 250401 (2003).

[13] C.A. Regal, M. Greiner and D.S. Jin, *Phys. Rev. Lett.* **92**, 040403 (2004).

[14] M. W. Zwierlein, A. Schirotzek, C.H. Schunck, and W. Ketterle, *Science* **311**, 492 (2006).

[15] R.A. Duine and A.H. MacDonald, *Phys. Rev. Lett.* **95**, 230403 (2005).

[16] "The Many-Body Challenge Problem" formulated by G. F. Bertsch, 1999, see R. F. Bishop, *Int. J. Mod. Phys. B* **15**, iii (2001).

[17] K. M. O'Hara et al., *Science* **298**, 2179 (2002);

[18] J. Kinast, A. Turlapov, J. E. Thomas, Q. Chen, J. Stajic, and K. Levin, *Science* **307**, 1296 (2005);

[19] T.-L. Ho, *Phys. Rev. Lett.* **92**, 090402 (2004);

[20] A. Bulgac, J.E. Drut, and P. Magierski, *Phys. Rev. Lett.* **96**, 090404 (2006).

[21] F. Chevy, "Universal phase diagram of a strongly interacting Fermi gas with unbalanced spin populations", *cond-mat/0605751* (2006).

[22] T. Marujama and G.F. Bertsch, *Phys. Rev. A* **73**, 013610 (2006).

[23] T. Sogo and H. Yabu, *Phys. Rev. A* **66**, 043611 (2002).

[24] L. Salasnich, B. Pozzi, A. Parola and L. Reatto, *J. Phys. B: At. Mol. Opt. Phys.* **33**, 3943 (2000).

[25] M. Brack and R. K. Bhaduri: *Semiclassical Physics* (Westview Press, Boulder, USA, 2003).

[26] P. Ring and P. Schuck, *The Nuclear Many-Body Problem*, (Springer-Verlag, New York, 1980).

[27] M. Gaudin, *Phys. Lett.* **24A**, 55 (1967).

[28] H. Hui, L. Xia-Ji and P.D. Drummond, "FFLO phase in one-dimensional polarized Fermi gases", *cond-mat/0610448*.

[29] B. P. van Zyl, R.K. Bhaduri, A. Suzuki, and M. Brack, *Phys. Rev. A* **67**, 023609 (2003).

[30] M. Ögren, unpublished (2004).

[31] Y. Yu, M. Ögren, S. Åberg, S.M. Reimann and M. Brack, *Phys. Rev. A* **72**, 051602(R) (2005).